

- Synopsis of subfamilies of Reduviidae
- Morphology
- Key to the subfamilies of Reduviidae
- Key to the tribes of Harpactorinae
- Key to the tribes of Holoptilinae
- Key to the tribes of Phymatinae
- Key to the tribes of Triatominae
- Key to the tribes of Tribelocephalinae
- Taxon treatments: subfamilies of Reduviidae
 - Introduction and general references
 - 1) Bactrodinae
 - 2) Centrocnemidinae
 - 3) Cetherinae
 - 4) Chryxinae
 - 5) Ectrichodiinae
 - 6) Elasmodeminae
 - 7) Emesinae
 - 8) Hammacerinae
 - 9) Harpactorinae
 - 10) Holoptilinae



Contents

- Taxon treatments (cont.)
 - 11) Manangocorinae
 - 12) Peiratinae
 - 13) Phimophorinae
 - 14) Phymatinae
 - 15) Physoderinae
 - 16) Pseudocetherinae
 - 17) Reduviinae
 - 18) Saicinae
 - 19) Salyavatinae
 - 20) Sphaeridopinae
 - 21) Stenopodainae
 - 22) Triatominae
 - 23) Tribelocephalinae
 - 24) Vesciinae
 - 25) Visayanocorinae

Synopsis of Subfamilies of Reduviidae

15 Physoderinae



16 Pseudocetherinae



17 Reduviinae



19 Salyavatinae



18 Saicinae



8 Hammacerinae



9 Harpactorinae



1 Bactroinae



2 Centrocnemidinae



3 Cetherinae



4 Chryxinae



10 Holoptiinae



11 Manangocorinae



20 Sphaeridopinae



21 Stenopodinae



12 Peiratinae



5 Ectrichodiinae



22 Triatominae



24 Vesciinae



13 Phimophorinae



6 Elasmodeminae



23 Tribelocephalinae



14 Phymatinae





7 Emesinae







25 Visayanocorinae







Morphology




1**1'**

1	Labium with 4 segments visible (separated by membrane)	<u>2</u>
1'	Labium with 3 segments visible (separated by membrane), segment 1 reduced	<u>3</u>





2 (1)	Pedicel subdivided into 4-36 pseudosegments, humeral angle of pronotum rounded <u>Hammacerinae</u> [Nearctic; Neotropical]	
2' (1)	Pedicel consisting of one uniform segment, humeral angle of pronotum with strong spines <u>Centrocnemidinae</u> [Oriental]	



3 (1) Cubitus (Cu) usually forming quadrate, pentagonal, or hexagonal cell; pedicel and flagellomeres folded underneath scape or claw with basal tooth (*the latter two characters are important should the specimen be apterous*) **4**



3' (1) Cubitus (Cu) without quadrate, pentagonal, or hexagonal cell; pedicel and flagellomeres not folded underneath scape or claw without basal tooth **5**



4 (3) Cubitus (Cu) usually forming pentagonal or hexagonal cell; if cubital cell absent, with prominent mandibular plate, pedicel and flagellomeres usually folded underneath scape, body often brown or pale.....**Stenopodainae** [Cosmopolitan]


4' (3) Cubitus (Cu) usually forming quadrate cell, mandibular plate small, pedicel and flagellomeres usually not folded under scape, body color variable**Harpactorinae** [Cosmopolitan]

See key to tribes




5 (3) Small (3-9 mm), membrane with a single cell, head short and transverse, anteriorly declivous, labium short and stout **Chryxinae** [Neotropical]

5' (3) Body size variable, membrane usually with 2 cells (rarely only 1) or 3 free veins, head short or elongate, rarely anteriorly declivous, labium usually elongate **6**

6fossula
spongiosa
absentfossula
spongiosa**6'**

tenent hairs



**6 (5)**

Apex of foretibia ventrally without obvious fossula spongiosa, i.e. never with specialized tenent hairs (acanthae with capitate apex)

7**6' (5)**


Apex of foretibia ventrally with obvious fossula spongiosa, i.e. an area densely beset with specialized tenent hairs (acanthae with capitate apex)

21




7 (6)	Vestiture on antenna and tibiae usually longer than twice the diameter of segment, often with trichome, foretibial comb absent; if setation dense but short, antenna with only 3 segments Holoptilinae [Circumtropical]	<u>See key to tribes</u>
7' (6)	Vestiture on antenna and tibiae, if present, usually shorter than diameter of segment, trichome absent, foretibial comb present; if antenna 3-segmented, then without dense setation	<u>8</u>


8





foretibial comb on spur





8'



foretibial comb not on spur




8 (7)	Foretibial comb located on prominent spur	<u>9</u>
8' (7)	Foretibial comb not located on prominent spur	<u>10</u>



9 (8) Body robust, third (second visible) labial segment (L3) straight, antenna short, scape and pedicel of about the same length, forecoxa short and rounded, claw symmetrical **Physoderinae** [Circumtropical]

9' (8) Body slender, third (second visible) labial segment (L3) gently curved, scape much longer than pedicel, forecoxa elongate, claw asymmetrical **Bactrodinae** [Neotropical]

10



L3 slender and straight

10'



L2

L3

L3 curved and usually slightly incrassate

1



L2

L3



10 (8)


Third (second visible) labial segment (L3) slender and straight
.....Triatominae (in part) [Cosmopolitan]

See key to tribes

10' (8)


Third (second visible) labial segment (L3) curved and usually slightly incrassate

11



membrane with 3 free ending veins, without cells

body in lateral view extremely flattened



body in lateral view usually not extremely flattened


if flattened, then membrane with 1-2 cells

11 (10) Body in lateral view extremely flattened, membrane with 3 free ending veins, without cells **Elasmodeminae** [Neotropical]

11' (10) Body in lateral view usually not extremely flattened; if flattened, then membrane with 1-2 cells

12

ocelli absent



woolly vestiture consisting of hairs that are bulbous basally and curved apically

pedicel + flagellomeres folded



ocelli present

12'



pedicel + flagellomeres not folded

glabrous or with sparse to dense vestiture, never with woolly vestiture



12 (11)



Body with woolly vestiture consisting of hairs that are bulbous basally and curved apically, ocelli absent, pedicel and flagellomeres often folded underneath scape **Tribelocephalinae** [Circumtropical]

See key to tribes

12' (11)

Body glabrous or with sparse to dense vestiture, never with woolly vestiture, ocelli present or absent, pedicel and flagellomeres not folded underneath scape


13



13 (12) Head with buccula obscuring the base of the labium **14**

13' (12) Head without buccula obscuring the base of the labium **15**

14



head almost twice as long as high

foreleg not chelate/sub-chelate



body flattened




14'



foreleg chelate

head about as long as high





forefemur with spines

body not flattened



foreleg subchelate





14 (13)

Foreleg not chelate or subchelate, forefemur without prominent spines, body somewhat flattened, head almost twice as long as highPhimophorinae [Neotropics, Oriental]

14' (13)


Foreleg usually chelate or subchelate; if not thus modified, then forefemur with prominent spines, body not flattened, head almost as high as longPhymatinae [Cosmopolitan]

See key to tribes



<p>15 (13)</p>	<p>Maxillary plate greatly enlarged, mandibular plate small, clypeus elongate and thus head appearing tri-lobed in dorsal view <u>Pseudocetherinae</u> [Circumtropical]</p>	
<p>15' (13)</p>	<p>Maxillary plate typically small, mandibular plate small or large, head not appearing tri-lobed</p>	<p><u>16</u></p>

16



spur

spur

spur

16'



foretibia without spur

16 (15)

Foretibia with spur


17

16' (15)

Foretibia without spur

18

17




L2 as long or slightly shorter than L3

scape as long as head



scape shorter than or as long as head

17'



L2 much longer than L3

scape much longer than head



17 (16) Second (first visible) labial segment (L2) as long or slightly shorter than third segment (L3), scape shorter than or as long as head
 **Vesciinae** [Neotropical]

17' (16) Second (first visible) labial segment (L2) much longer than third segment (L3), scape much longer than head
 **Visayanocorinae** [Paleotropical]

18

ocelli absent



ocelli present


18'




ocelli present

<p>18 (16)</p>	<p>Ocelli absent, even in macropterous forms; if ocelli present (1 genus of Emesinae) forecoxa usually at least 4 times as long as wide</p>	<p>19</p>
<p>18' (16)</p>	<p>Ocelli present</p>	<p>20</p>


19



19'




<p>19 (18)</p>	<p>Forecoxa usually at least 4 times as long as wide, acetabulum of foreleg opening anterior Emesinae [Cosmopolitan]</p>	
<p>19' (18)</p>	<p>Forecoxa at most 3 times as long as wide, acetabulum of foreleg opening ventrad; third visible labial segment (L3) often bulbous Saicinae [Cosmopolitan]</p>	



20

labium straight



large

5 mm



tarsi 3-segmented



20'

labium curved



small


5 mm



20 (18) Large, with small head, labium straight and slender, tarsi 3-segmented, color brown (or red and black in species with fossula spongiosa) **Sphaeridopinae (in part)** [Neotropical]

20' (18) Small, labium curved and stout, tarsi 2-segmented, wing venation and color as in Plate 20' **Manangocorinae** [Oriental]

21




transverse sulcus located behind middle of pronotum



forecoxa usually ~2 times as long as wide; flat laterally

21'



transverse sulcus at or in front of middle of pronotum



forecoxa usually less than 2 times as long as wide; rounded laterally

21 (6) Transverse sulcus on pronotum located behind middle of pronotum, forecoxa usually ~2 times as long as wide and with lateral surface flat **Peiratinae** [Cosmopolitan]

21' (6) Transverse sulcus on pronotum usually located at or in front of middle of pronotum, forecoxa usually less than 2 times as long as wide and without lateral surface flat **22**

22



L3 slender and straight

scutellum without posteriorly projecting prongs




22'



L3 curved and incrassate

if L3 slender and straight, then scutellum with posteriorly projecting prongs



22 (21)

Third (second visible) labial segment (L3) slender and straight; scutellum never with posteriorly projecting lateral prongs
..... Triatominae (in part) [Cosmopolitan]

See key to tribes

22' (21)

Second visible (third) labial segment (L3) curved and usually slightly incrassate; if slender and straight, then with two posteriorly projecting lateral prongs

23


23

scutellum with two prongs and without median tip



scape


pedicel



subdivided flagellomeres

23'


scutellum with one median tip



pedicel

scape

flagellomeres not subdivided



23 (22)

Scutellum usually with two posteriorly projecting lateral prongs and usually without the median tip, flagellomeres usually subdivided into 4-6 pseudosegments **Ectrichodiinae** [Cosmopolitan]


23' (22)

Scutellum with one median tip and without posteriorly projecting lateral prongs, flagellomeres not subdivided into pseudosegments

24

24

head elongate



if head short, then eyes usually reniform



24'

semi-globular eyes medium-sized, slightly to strongly pedunculate



head short

semi-globular eyes large, almost covering lateral head surface





24 (23)

Head usually elongate; if short, then eyes usually reniform and neither strongly pedunculate nor covering almost the entire lateral surface of the head **Reduviinae** [Cosmopolitan]

24' (23)

Head short, with semi-globular eyes that are either medium-sized and slightly to strongly pedunculate or large and almost covering the entire lateral surface of the head

25




25 (24) Tarsal formula 2-3-3, habitus as in Plate 25, sometimes with median, spine-like frontal process or dilated foretibia
 **Salyavatinae** [Circumtropical]

25' (24) Tarsal formula 3-3-3, habitus as in Plate 25', never with dilated foretibia; if frontal process present, then process bi-lobed

26

eyes pedunculate



labium stout and curved

brownish

26'

eyes not pedunculate



black and red



labium slender and straight



26 (25)

Eyes pedunculate, labium stout and curved, body mottled brownish

.....Cetherinae [Afrotropical; Neotropical]

26' (25)

Eyes not pedunculate, labium slender and straight, body black and red



.....Sphaeridopinae (in part) [Neotropical]

Key to the Tribes of Harpactorinae





To Tribe [Key](#)

Back to Subfamily [Key](#)



1	Labium curved Harpactorini [Cosmopolitan]	
1'	Labium straight	<u>2</u>



2 (1)	Clypeus enlarged into laterally flattened projection Dictyoteliini [Oriental]	
2' (1)	Clypeus short and rounded, not enlarged into projection	3

3



body elongate, appendages slender



3'





body robust, appendages not very slender





3 (2) Body elongate, appendages very slender and long
..... **Rhaphidosomini** [Afrotropical; Oriental]

3' (2) Body robust, appendages not very slender and long



4



<p>4 (3)</p>	<p>Labium extremely slender, reaching to middle of forecoxa Tegeini [Afrotropical; Oriental]</p>	
<p>4' (3)</p>	<p>Labium short and usually moderately slender or stout; if slender, then at most reaching anterior margin of forecoxa</p>	<p>5</p>



5 (4)	Pronotum completely covering scutellum Diaspidiini [Afrotropical]	
5' (4)	Scutellum at least in part exposed	6



<p>6 (5)</p>	<p>Foretarsus 1-segmented, habitus as in Plate 6 Ectinoderini [Oriental]</p>	
<p>6' (5)</p>	<p>Foretarsus 2-segmented, habitus as in Plate 6' Apiomerini [Neotropical]</p>	

Key to the Tribes of Holoptilinae





Modified from
Wygodzinsky and
Usinger (1963)




To Tribe [Key](#)

Back to Subfamily [Key](#)



1	Setae of antenna and legs shorter than diameter of respective segments, trichome absent Aradellini [Australia]	
1'	Setae of antenna and legs longer than diameter of respective segments, trichome absent or present	<u>2</u>

2




2 or 3 longitudinal veins not connected beyond middle of membrane

2'



3 longitudinal veins, inner and middle veins joining subapically to form elongate cells



trichome present



<p>2 (1)</p>	<p>Membrane with 2 or 3 free longitudinal veins not connected beyond middle of membrane; trichome absent; usually 5 mm or less Dasycnemini [Circumtropical]</p>
<p>2' (1)</p>	<p>Membrane with 3 longitudinal veins, at least inner and middle veins joining subapically to form an elongate cell; trichome present; usually 5 mm or longer Holoptilini [Afrotropical; Oriental; Australian]</p>

Key to the Tribes of Phymatinae




To Tribe [Key](#)

Back to Subfamily [Key](#)




1	Forefemur and foretibia forming a chelate or subchelate raptorial leg, forefemur ventrally not with strong spines	<u>2</u>
1'	Forefemur and foretibia not forming a chelate or subchelate raptorial leg, forefemur ventrally with strong spines Themonocorini [Afrotropical]	



2



2'



2 (1)	Foreleg subchelate (i.e., tibia clasped against enlarged body of femur)	<u>3</u>
2' (1)	Foreleg chelate (i.e., tibia clasped against distal process on femur) Carcinocorini [Oriental]	



3 (<u>2</u>)	Scutellum small and triangular Phymatini [Cosmopolitan]	
3' (<u>2</u>)	Scutellum large and tongue-shaped Macrocephalini [Cosmopolitan]	



Key to the Tribes of Triatominae

Modified from Lent and Wygodzinsky (1979)



To Tribe [Key](#)

Back to Subfamily [Key](#)





1 Ocelli not elevated, situated at level of integument, inconspicuous among coarse head granules or in or very close to interocular sulcus

2



1' Ocelli situated on distinct elevations on disc of postocular portion of head

3





2 (1) Head usually elongate, not strongly convex dorsally; maxillary plate projecting beyond apex of clypeus; interocular sulcus obsolete; corium with veins distinct; integument heavily granulose... **Bolboderini** [Neotropical]

2' (1) Head ovoid, strongly convex dorsally; maxillary plate small; interocular sulcus strongly backwardly curved; corium with veins obsolete; integument smooth, hairy **Cavernicolini** [Neotropical]



3 (1) Head elongate, insertion of antenna close to apex of head
**Rhodniini** [Neotropical]

3' (1) Head elongate or short and wide, if elongate then antenna inserted remote from apex of head **4**



<p>4 (3)</p>	<p>Body length ~ 5 mm; head very short and wide; hemelytra with small branch connecting basal portion of R+M to Sc Alberproseniini [Neotropical]</p>	
<p>4' (3)</p>	<p>Body length > 5 mm; head elongate; small branch connecting basal portion of R+M to Sc absent Triatomini [Circumtropical]</p>	



Key to the Tribes of Tribelocephalinae

Modified from Maldonado (1996)



To Tribe [Key](#)

Back to Subfamily [Key](#)



1	Apterous, eyes absent, tarsi 1-segmented..... Xenocaucini [Afrotropical]	
1'	Winged, with eyes, tarsi 2- or 3-segmented	2

2

m-cu cross vein

base of inner discal cell



2'

m-cu cross vein absent

base of inner discal cell



2 (1) Hemelytron with base of inner discal cell divided in two by cross vein m-cu **Opistoplatyini** [Afrotropical, Oriental, Australian]

2' (1) Hemelytron with base of inner discal cell not divided in two by a cross vein **Tribelocephalini** [Afrotropical, Oriental, Australian]

Taxon Treatments: Subfamilies of Reduviidae

Triatoma dispar, Costa Rica



© Stephen Marshall

Acanthaspis sulcipes, Ivory Coast



© Jean-Michel Bérénger

Agriocoris flavipes, Ecuador



© Stephen Marshall

Triatoma sanguisuga, United States



© Stephen Marshall

Taxon treatments: The following taxon treatments are designed to combine short diagnoses, summaries of the taxonomic history and distribution of the subfamily, some notes on the natural history, the most important references for each of the subfamilies, and additional images of specimens, dead and, where available, alive and in their natural surroundings. Diagnoses of the 25 subfamilies of Reduviidae here recognized are meant to complement the identification key. The taxonomic scope and history varies in depth from a fairly comprehensive treatment for some of the smaller subfamilies, to a rather superficial perspective for some of the larger and more diverse groups. The taxonomic notes are complemented by comments on the phylogenetic status of many of the subfamilies, making it evident that the classification of Reduviidae we accepted for the purposes of this project will be in flux in the near future. To shorten the list of references on each taxon treatment, we list below several references that are of importance to all of the taxon pages and that we do not repeat for individual subfamilies. The hyperlinks “Back to Content” and “Back to Key” point back to the main content page and the final page in the subfamily key for a given taxon, respectively.

Anatomical orientation: When referring to specific surfaces of appendages, in particular the legs, we think of the appendages being extended. For example, we refer to surface of the foretibia that carries the foretibial comb as anterior surface.

General references

- Henry, T.J. and Froeschner, R.C. 1988.** Catalog of the Heteroptera, or true bugs, of Canada and the Continental United States. CRC Press, Boca Raton.
- Hwang, W.S. and Weirauch, C. 2012.** Evolutionary history of assassin bugs: insights from divergence dating and ancestral state reconstruction. PLoS ONE, **7**: 1– 12. e45523. doi:10.1371/journal.pone.0045523
- Maldonado, J. 1990.** Systematic catalogue of the Reduviidae of the world (Insecta: Heteroptera). Caribbean Journal of Science, Special Edition.
- Maw, H.E.L., Foottit, R.G., Hamilton, K.G.A., and Scudder, G.G.E. 2000.** Checklist of the Hemiptera of Canada and Alaska. NRC Research Press, Ottawa.
- Putchkov, V.G. and Putchkov, P.V. 1985.** A catalog of assassin-bug genera of the World (Heteroptera, Reduviidae). [Published by the authors], Kiev.
- Schuh, R.T. and Slater, J.A. 1995.** True bugs of the World (Hemiptera: Heteroptera): classification and natural history. Cornell University Press, Ithaca.

Bactrodinae


[Back to Content](#) [Back to Key](#)

Diagnosis: Bactrodinae are characterized by the slender body, gently curved third (second visible) labial segment, elongated postocular region, elongate forecoxa, asymmetrical claws, scape longer than the pedicel, foretibial comb on prominent spur, single cell in the forewing membrane, bifurcated median pygophore process, and the ventral spine on the foretrochanters. The third tarsal segment on the mid- and hindlegs is incrassate and has a stout spine near the middle of the ventral surface.

Taxonomic Notes and Distribution: This subfamily comprises 5 Neotropical species in the genus *Bactrodes* Stål that are distributed from Mexico to Argentina (Coscarón and Melo 2003). McAtee and Malloch (1923) and Coscarón and Melo (2003) provided keys to species. Bactrodinae were first treated as a subfamily by Stål (1862). Davis (1969) considered Bactrodinae to be closely related to, but distinct from, the Harpactorinae. Recent phylogenetic analyses have failed to include Bactrodinae, and the phylogenetic position of this subfamily remains to be investigated.

Occurrence in Canada: Bactrodinae do not occur in Canada.

Natural History: Bérenger and Pluot-Sigwalt (1997) reported *Bactrodes femoratus* (Fabricius) in French Guiana to be associated with plants in the family Melastomataceae.



[Next](#)

Bactrodinae (cont.)



Bactrodes femoratus, Trinidad

© Christiane Weirauch



Bactrodes sp., Colombia

© Guanyang Zhang

References

- Bérenger, J. and Pluot-Sigwalt, D. 1997.** Special relationships of certain predatory Heteroptera Reduviidae with plants. First known case of a phytophagous Harpactorinae. *Comptes Rendus de l'Academie des Sciences Series III Sciences de la Vie*, **320**: 1007-1012.
- Coscarón, M.C. and Melo, M.C. 2003.** Revision of the subfamily Bactrodinae (Heteroptera, Reduviidae), with a phylogenetic analysis of *Bactrodes*. *Zootaxa*, **304**: 1-15.
- Davis, N.T. 1969.** Contribution to the morphology and phylogeny of the Reduvidae. Part IV. The harpactoroid complex. *Annals of the Entomological Society of America*, **62**: 74-94.
- McAtee, W.L. and Malloch, J.R. 1923.** Notes on American Bactrodinae and Saicinae (Heteroptera: Reduviidae). *Annals of the Entomological Society of America*, **16**: 247-254.
- Stål, C. 1862.** Hemiptera Mexicana enumeravit speciesque novas descripsit. *Stettiner Entomologische Zeitung*, **23**: 437-462.

Centrocnemidinae

[Back to Content](#)

[Back to Key](#)

Diagnosis: Centrocnemidinae are diagnosed by 4 visible labial segments, the pedicel consisting of one uniformly sclerotized segment, strong spines on the humeral angle of the pronotum, 2 elongate cells in the forewing membrane, and the tuberculate body.

Taxonomic Notes and Distribution: About 33 extant species in 4 genera are currently recognized in this subfamily (Miller 1956; Truong et al. 2010; Rédei and Tsai 2011). See Rédei and Tsai (2011) for a discussion on the recent change of the subfamily name. All recent taxa are Oriental in distribution (Maldonado 1990), but reduviid fossils that may belong to this subfamily are known from Baltic Amber (Popov and Putschkov 1998). Comparative morphology and phylogenetic analyses indicate that Centrocnemidinae are part of the phymatine complex of reduviid subfamilies (Clayton 1990; Weirauch et al. 2011; Hwang and Weirauch 2012).

Occurrence in Canada: This subfamily does not occur in Canada.

Natural History: Most species are collected on tree trunks (Schuh and Slater 1995).

[Next](#)

Neocentrocnemis signoreti, Singapore



Centrocnemidinae sp., Singapore



Centrocnemidinae (cont.)

[Previous](#)

References

- Clayton, R.A. 1990.** A phylogenetic analysis of the Reduviidae (Hemiptera: Heteroptera) with redescription of the subfamilies and tribes. Ph.D. thesis, The George Washington University, Washington, DC.
- Miller, N.C.E. 1956.** Centrocneminae, a new sub-family of the Reduviidae (Hemiptera-Heteroptera). Bulletin of the British Museum (Natural History), Entomology, **4**: 219-283.
- Popov, Y. and Putschkov, P.V. 1998.** *Redubinotus liedtkei* n. gen. n. sp. - the second Centrocnemina from the Baltic amber (Heteroptera: Reduviidae: Centrocnemidinae). Annals of the Upper Silesian Museum in Bytom Entomology, **8-9**: 205-210.
- Rédei, D. and Tsai, J.F. 2011.** The assassin bug subfamilies Centrocnemidinae and Holoptilinae in Taiwan (Hemiptera: Heteroptera: Reduviidae). Acta Entomologica Musei Nationalis Pragae, **51**: 411-442.
- Truong, X.L., Li, H., and Cai, W. 2010.** First record of the assassin bug subfamily Centrocnemidinae (Hemiptera: Heteroptera: Reduviidae) from Vietnam, with the description of a new species. Zootaxa, **2347**: 64-68.
- Weirauch, C., Forero, D., and Jacobs, D.H. 2011.** On the evolution of raptorial legs – an insect example (Hemiptera: Reduviidae: Phymatinae). Cladistics, **27**:138-149.



Centrocnemidinae sp.,
Singapore

Cetherinae

[Back to Content](#)

Diagnosis: Cetherinae are diagnosed by the pedunculate compound eyes, stout and curved labium, presence of ocelli and fossula spongiosa, and the transverse sulcus on the vertex.

Taxonomic Notes and Distribution: About 23 species in 5 genera are currently classified in this subfamily. Villiers (1948) revised the 4 Afrotropical genera; the Neotropical genus *Eupheno* Gistel comprises only 3 described species. Based on a recent molecular phylogenetic analysis of Reduviidae that included Old and New World representatives of Cetherinae, this group may be polyphyletic (Hwang and Weirauch 2012). Cetherinae were first treated as a tribe within the Reduviinae by Jeannel (1919) and the group was given subfamily status by Miller (1955; as Eupheninae). These authors recognized 3 tribes, the Cetherini, Euphenini, and Pseudocetherini. The latter is here treated as a distinct subfamily (in agreement with Maldonado [1990]). Pending a combined phylogenetic analysis of Reduviidae, Cetherini and Euphenini may in the future be treated as separate subfamilies.

Occurrence in Canada: This subfamily does not occur in Canada.

Natural History: Some species are known to prey on termites (Miller 1956). Species of *Eupheno* and *Cethera* Amyot and Serville are found on the underside of fallen tree trunks and logs and similar hidden microhabitats that are typically also inhabited by termites.

[Back to Key](#)

Cethera musiva,
Madagascar



Eupheno sp.,
Peru



© Christiane Weirauch

[Next](#)

Cetherinae (cont.)

[Previous](#)



References

- Hwang, W.S. and Weirauch, C. 2012.** Evolutionary history of assassin bugs: insights from divergence dating and ancestral state reconstruction. *PLoS ONE*, **7**(9): 1-12. e45523. doi:10.1371/journal.pone.0045523.
- Jeannel, R. 1919.** Voyage de Ch. Alluaud et R. Jeannel en Afrique Orientale (1911-1912). Resultats scientifiques. Insectes Hemipteres. III. Hencocephalidae et Reduviidae. L. Lhomme, Paris. pp. 133-313.
- Miller, N.C.E. 1955.** The synonymy of *Eupheno* Gistel (Hemiptera-Heteroptera, Reduviidae, Reduviinae) and the description of a new subfamily. *Annals and Magazines of Natural History*, **8**: 449-452.
- Miller, N.C.E. 1956.** *The Biology of the Heteroptera*. Leonard Hill Books, London.
- Villiers, A. 1948.** Faune de l'Empire Francais. IX. Hemipteres Reduviides de l'Afrique Noire. Office de la Recherche Scientifique Coloniale, Editions du Museum, Paris.

Chryxinae

Diagnosis: Chryxinae are small (3-9 mm) assassin bugs with a short, stout labium with 3 visible segments; a short, stout, transverse, anteriorly declivous head; and a single cell in the forewing membrane.

Taxonomic Notes and Distribution: This subfamily comprises 5 Neotropical species in 4 genera (Gil-Santana et al. 2007; Weirauch 2012). Generic limits within the subfamily may be doubtful (Weirauch 2012). Champion (1897-1901) first described this group by giving subfamily status to the genus *Chryxus* Champion. Chryxinae are restricted to the Neotropics.

Occurrence in Canada: Chryxinae do not occur in Canada.

Natural History: The biology of species in this subfamily is unknown. *Petasolentia goellnerae* Weirauch was collected at light in the Andean foothills of Southeastern Peru.

References

- Champion, G.C. 1897-1901.** Biologia Centrali Americana. Insecta. Rhynchota. Hemiptera-Heteroptera. Vol. II.
- Gil-Santana, H.R., Costa, L.A.A., and Marques, O.M. 2007.** Sinopse dos Chryxinae (Hemiptera, Reduviidae). Revista Brasileira de Zoologia, **24**: 77-83.
- Weirauch, C. 2012.** *Petasolentia*, a new genus and species of Chryxinae (Hemiptera: Reduviidae). Entomologische Zeitschrift, **122**: 119-122.

[Back to Content](#) [Back to Key](#)



Chryxus tomentosus,
Panama

© Christiane Weirauch



Petasolentia goellnerae,
Peru

Ectrichodiinae


[Back to Content](#)

[Back to Key](#)

Diagnosis: The Ectrichodiinae, or millipede assassin bugs, are typically diagnosed by the presence of a bifurcated scutellum (2 lateral posteriorly projecting processes), subdivided antennal basi- and distiflagellomeres (flagellum appears 4- to 6-segmented), forewing membrane with 2 or 3 cells, and the fossula spongiosa on the fore- and midtibiae. There are several taxa that do not possess all of these diagnostic features, such as *Zirta* Stål and *Schuhella* Dougherty (flagellomeres not subdivided) and some species of *Katanga* Schouteden and *Cimbus* Hahn (without fossula spongiosa). The lateral processes of the scutellum are very short in some taxa, especially in apterous forms.

Taxonomic Notes and Distribution: More than 660 species in ~120 genera have been described (Maldonado 1990; Dougherty 1995; Carpintero and Maldonado 1996; Bérenger and Gil-Santana 2005; Weirauch et al. 2009; Chlond 2010). More than half of the described genera are monotypic and no tribes are recognized. Amyot and Serville (1843) described Ectrichodiinae as a subfamily. Dougherty (1995) provided a review of the taxonomic history for the subfamily. This subfamily is largely restricted to the tropical and subtropical areas around the world, but some taxa also occur in temperate regions.

[Next](#)



Ectrichodiinae (cont.)

Occurrence in Canada: Ectrichodiinae have not been recorded from Canada.

Natural History: Many, if not all, Ectrichodiinae are in all likelihood specialized millipede predators (Forthman and Weirauch 2012). Aposematic or metallic coloration and strong sexual dimorphism are common in this group. Some species in the Neotropical and Afrotropical regions exhibit wing polymorphism within species (Dougherty 1995). Ectrichodiinae are frequently found in leaf litter and are predominantly nocturnal (and often attracted to artificial lights), although some species may also be active during the day (Miller 1953; Louis 1974; Schuh and Slater 1995).

Glymmatophora sp., Namibia



© Stephen Marshall

Ectrichodiinae sp.,
Singapore



© Melvyn Yeo

Neoscadra sp., Singapore



© Melvyn Yeo

References

Amyot, C.J.B. and Serville, A. 1843. Histoire Naturelle des Insectes. Hemipteres. Roret, Paris.

Bérenger, J.M. and Gil-Santana, H. 2005. Nouveau genre et nouvelles espèces d'Ectrichodiinae d'Amérique Du Sud (Heteroptera, Reduviidae). Bulletin de la Société entomologique de France, **110**: 509-516.

Carpintero, D.J. and Maldonado, J. 1996. Diagnostic characters and key to the genera of American Ectrichodiinae (Heteroptera, Reduviidae). Caribbean Journal of Science, **32**: 125-141.

Ectrichodiinae (cont.)

[Previous](#)

References (cont.)

- Chlond, D. 2010.** A new, remarkable genus and two new species of Ectrichodiinae (Hemiptera: Heteroptera: Reduviidae) from Madagascar. *Zootaxa*, **2522**: 61-68.
- Dougherty, V. 1995.** A review of the New World Ectrichodiinae genera (Hemiptera: Reduviidae). *Transactions of the American Entomological Society*, **121**: 173-225.
- Forthman, M. and Weirauch, C. 2012.** Toxic associations: a review of the predatory behaviors of millipede assassin bugs (Hemiptera: Reduviidae: Ectrichodiinae). *European Journal of Entomology*, **109**: 147-153.
- Louis, D. 1974.** Biology of Reduviidae of cocoa farms in Ghana. *American Midland Naturalist*, **91**: 68-89.
- Miller, N.C.E. 1953.** Notes on the biology of the Reduviidae of southern Rhodesia. *Transactions of the Zoological Society of London*, **27**: 541-672.
- Weirauch, C., Rabitsch, W., and Rédei, D. 2009.** *Austrokatanga*, gen. nov., new genus of Ectrichodiinae (Hemiptera: Heteroptera: Reduviidae) from Australia. *Zootaxa*, **2094**: 1-15.



© Melvyn Yeo

Schottus sp., Thailand



Ectrychotes sp., Vietnam

© Stephen Marshall

Elasmodeminae

[Back to Content](#)

[Back to Key](#)

Diagnosis: The subfamily Elasmodeminae is distinguished by the dorsoventrally flattened body and the 3 free veins reaching the posterior margin of the forewing membrane. Other features that characterize this subfamily include the presence of a fossula spongiosa on the foretibia, 3 visible labial segments, and the spiny legs.

Taxonomic Notes and Distribution: This subfamily is comprised of 3 Neotropical species (southern Brazil, Paraguay, and Argentina) in the genus *Elasmodema* Stål. Several authors have treated Elasmodeminae as a distinct family (Wygodzinsky 1944; China and Miller 1959; Maldonado 1990), but Carayon et al. (1958) convincingly argued for its inclusion into the Reduviidae, based on a close relationship with the Holoptilinae.

Occurrence in Canada: Elasmodeminae do not occur in Canada.

Natural History: Species live under bark and prey on other insects (Wygodzinsky 1944).

References

Carayon, J., Usinger, R.L., and Wygodzinsky, P.W. 1958. Notes on the higher classification of the Reduviidae, with the description of a new tribe of the Phymatinae (Hemiptera- Heteroptera). *Revue de zoologie et de botanique africaine*, **57**: 256-281.

China, W.E. and Miller, N.C.E. 1959. Checklist and keys to the families and subfamilies of the Hemiptera-Heteroptera. *Bulletin of the British Museum (Natural History) (Entomology)*, **8**: 1-45.

Wygodzinsky, P.W. 1944. Contribuição ao conhecimento do genero *Elasmodema* Stål, 1960 (Elasmodemidae, Reduvioidea, Hemiptera). *Revista Brasileira de Biologia*, **4**: 193-213.



Emesinae

[Back to Content](#) [Back to Key](#)

Diagnosis: Emesinae, or thread-legged bugs, are characterized by the elongate and slender body, typically thread-like legs with the forecoxa usually at least 4 times as long as wide, acetabulum of the foreleg opening anteriorly, absence of ocelli (except in *Armstrongocoris* Wygodzinsky) and the tibial fossula spongiosa, and the often asymmetrical pretarsal claws of the forelegs.

Taxonomic Notes and Distribution: Wygodzinsky (1966) produced a monographic revision of the extant species in the subfamily and recognized 6 tribes: Collartidini, Leistarchini, Emesini, Ploiariini, Deliastini, and Metapterini. The group comprises >900 extant species in >90 genera (Maldonado 1990) and a number of fossil taxa (see Popov et al. 2011). Wygodzinsky considered the Emesinae to be closely related with the Saicinae and Visayanocorinae, a scenario that is supported by recent phylogenetic analyses (Weirauch 2008; Weirauch and Munro 2009; Hwang and Weirauch 2012). Emesinae are cosmopolitan, but diversity is highest in the tropics. The group shows significant species diversity on Pacific islands, areas not typically inhabited by a diverse fauna of Reduviidae.



Emesinae (cont.)

[Previous](#)

[Next](#)

Occurrence in Canada: Species in the genera *Barce* Stål, *Emesaya* McAtee and Malloch, and *Empicoris* Wolff occur in Canada.

Natural History: Emesinae are cryptic and often found associated with dense vegetation, tree trunks, leaf litter, and spider webs (Wygodzinsky 1966). Some species appear to be obligate predators of web-building spiders, and some are facultatively associated with spiders and spider prey (Gillet 1957; Santiago-Blay and Maldonado 1988; Forero 2007; Wignall and Taylor 2008). A number of species of Emesinae are known to be cave-dwelling (Gagne and Howarth 1974; Maldonado 1994; Ribes et al. 1997; Pape 2013), a microhabitat that harbors otherwise only few reduviid species.

Emesinae sp., Bolivia



© Stephen Marshall

Empicoris sp., Canada



© Stephen Marshall

© Melvyn Yeo

References

- Forero, D. 2007.** Description of a new species of *Liaghinella* (Hemiptera: Heteroptera: Reduviidae: Emesinae) from the Colombian Andes, with notes on its feeding habits and conservation status. *Zootaxa*, **1502**: 55-68
- Gagne, W.C. and Howarth, F.G. 1974.** The cavernicolous fauna of Hawaiian lava tubes, Part VII. Emesinae or thread-legged bugs (Heteroptera: Reduviidae). *Pacific Insects*, **16**: 415-426.
- Gillet, J.D. 1957.** On the habits and life history of captive emesine bugs (Hemiptera: Reduviidae). *Proceedings of the Royal Entomological Society London Series A*, 32, **10-12**: 193-195.

Emesinae (cont.)

[Previous](#)

Reference (cont.)

- Maldonado 1994.** Hemiptera-Heteroptera. *In* Encyclopaedia Bioespeologica. *Edited by* Juberthie and Decu. Bucarest, Rumania. Tomo I. Pp. i-xii, 7-311.
- Pape, R.B. 2013.** Description and ecology of a new cavernicolous, arachnophilous thread-legged bug (Hemiptera: Reduviidae: Emesini) from Kartchner Caverns, Cochise County, Arizona. *Zootaxa*, **3670**: 137-156.
- Popov, Y. A., Kosmowska-Ceranowicz, B., Herczek, A., and Kupryjanowicz, J. 2011.** Review of true bugs (Insecta: Hemiptera, Heteroptera) from the amber collection of the Museum of the Earth of PAS in Warsaw with some remarks on heteropteran insects from Eocene European amber. *Polskie Pismo Entomologiczne, Gydnia*, **80**: 699-728.
- Ribes R., Oromi P., and Ribes E. 1997.** Una nueva *Collartida* Villiers, 1949 subterranea de La Palma, islas Canarias (Heteroptera, Reduviidae, Emesinae). *Bulletin de la Société entomologique de France*, **26**: 99-105.
- Santiago-Blay, J.A. and Maldonado, J. 1988.** Observations on the true bugs *Emesa tenerrima*, a possible spider mimic and *Ghilianella borincana* (Hemiptera: Reduviidae: Emesinae) from Puerto Rico. *Proceedings of the Entomological Society of Washington*, **90**: 369-372.
- Weirauch, C. 2008.** Cladistic analysis of Reduviidae (Heteroptera: Cimicomorpha) based on morphological characters. *Systematic Entomology*, **33**: 229-274.
- Weirauch, C. and Munro, J.B. 2009.** Molecular phylogeny of the assassin bugs (Hemiptera: Reduviidae), based on mitochondrial and nuclear ribosomal genes. *Molecular Phylogenetics and Evolution*, **53**: 287-299.
- Wignall, A.E. and Taylor, P.W. 2008.** Biology and life history of the araneophagic assassin bug *Stenolemus bituberus* including a morphometric analysis of the instars (Heteroptera, Reduviidae). *The Journal of Natural History*, **42**: 59-76.
- Wygodzinsky, P.W. 1966.** A monograph of the Emesinae (Reduviidae, Hemiptera). *Bulletin of the American Museum of Natural History*, **133**: 1-614.

Hammacerinae

Diagnosis: Hammacerinae are characterized by the 4-segmented labium, the antennal pedicel being subdivided into 4-36 pseudosegments, the head strongly produced anteriorly and not constricted or elongated behind the compound eyes, the forewing membrane with 2 closed cells, and the fossula spongiosa on fore- and midlegs.

Taxonomic Notes and Distribution: This group was originally proposed as a subfamily by Stål (1859), and is sometimes referred to as Microtominae. The subfamily consists of 2 genera and ~19 described species that occur in the New World: *Homalocoris* Perty and *Microtomus* Illiger. *Microtomus* was revised by Stichel (1926), with subsequent taxonomic publications focusing on species occurring in South America (e.g., Coscarón and Giacchi 1985; Melo and Coscarón 2004). Maldonado (1987, 1991) treated *Homalocoris*.

Occurrence in Canada: This subfamily is not known to occur in Canada.

Natural History: Hammacerinae are typically found underneath bark (Radio 1927).

[Back to Content](#)

[Back to Key](#)

Microtomus sp., Colombia



© Dimitri Forero

[Next](#)

Hammacerinae (cont.)

References

- Coscarón, M.C. and Giacchi, J.C. 1985.** Revision de la subfamilia Microtominae (Hemiptera, Reduviidae). I. *Microtomus lunifer* (Berg). Revista de la Sociedad Entomológica Argentina, **44**: 243-250.
- Maldonado, J. 1987.** *Homalocoris punctatus* n. sp. and key to the species in the genus (Reduviidae, Microtominae). Journal of Agriculture of the University of Puerto Rico, **71**: 249-253.
- Maldonado, J. 1991.** Classification of *Homalocoris* (Heteroptera: Reduviidae: Hammacerinae), with the description of a new species. Proceedings of the Entomological Society of Washington, **93**: 703-708.
- Melo, M.C. and Coscarón, M.C. 2004.** Redescription of *Microtomus reuteri* Berg (Heteroptera: Reduviidae: Hammacerinae) from Southern South America. Entomological News, **115**: 249-254.
- Radio, P.A. 1927.** Studies on the biology of the Reduviidae of American north of Mexico. The University of Kansas science bulletin, **17**: 5-291.
- Stål, C. 1859.** Nova methodus Reduvina (Burm.) disponendi. Berliner Entomologische Zeitschrift, **3**: 328.
- Stichel, W. 1926.** Die Gattung *Microtomus* Illiger (Hem., Het., Reduv.). Deutsche Entomologische Zeitschrift, **1926**: 179-190.



Homalocoris sp., Brazil

Harpactorinae

[Back to Content](#) [Back to Key](#)

Diagnosis: Harpactorinae are characterized by the presence of a quadrate cell on the corium formed by the cubitus and a well-developed subapical spur on the foretibia that carries the foretibial comb. Most Harpactorinae also have a cylindrical postocular region and an elongated antennal scape.

Taxonomic Notes and Distribution: With more than 2,800 described extant species in ~320 genera, Harpactorinae is the largest subfamily of Reduviidae. Harpactorinae have been recognized as a group since Amyot and Serville (1843), but the tribal classification has been in flux, with Davis (1969) and Schuh and Slater (1995) recognizing 6 tribes (Apiomerini, Diaspidiini, Ectinoderini, Harpactorini, Rhaphidosomini, and Tegeini), but others also accepting Dicrotelini as a separate tribe (Miller 1954; Tomokuni and Cai 2002). We follow the latter and recognize 7 tribes in the subfamily. Members of this group are variable in body shape and color, with many species being rather colorful. Among the most visually striking Harpactorinae are the resin bugs (Apiomerini, Ectinoderini, and Diaspidiini; Forero et al. 2013), the wasp or bee mimicking bugs (e.g., *Hiranetis* Spinola, *Notocyrtus* Burmeister, and *Coilopus* Elkins; Zhang and Weirauch 2013a), the wheel bugs in the New World genus *Arilus* Hahn, stick-like Rhaphidosomini, and species of the Oriental *Eulyes* Amyot and Serville and related taxa.



[Next](#)

Harpactorinae (cont.)

[Previous](#)

[Next](#)

Taxonomic Notes and Distribution (cont.): Phylogenetic analyses support the monophyly of Harpactorinae, but relationships among tribal-level taxa within the group are in need of additional research: Harpactorini were recently shown to be paraphyletic or even polyphyletic (Hwang and Weirauch 2012; Forero et al. 2013; Zhang and Weirauch 2013a). Harpactorinae and the tribe Harpactorini have a worldwide distribution; Apiomerini are restricted to the New World; Diaspidiini and Rhaphidosomini are Afrotropical; Ectinoderini are restricted to the Oriental region; and Dicrotelini and Tegeini occur in the Old World tropics.


Occurrence in Canada: Genera recorded from Canada are *Acholla* Stål, *Apiomerus* Hahn, *Arilus* Hahn, *Fitchia* Stål, *Pselliopus* Bergroth, *Rhynocoris* Hahn, *Sinea* Amyot and Serville, and *Zelus* Fabricius.



Harpactorinae (cont.)

[Previous](#)[Next](#)

Natural History: The biology of Harpactorinae shows tremendous variation. The majority of species are diurnal and frequently found on plants, but some species are ground-dwelling or bark-inhabiting. Some harpactorine species show close association with specific plants (Bérenger and Pluot-Sigwalt 1997). Due to the general predacious habits of many species, some Harpactorinae are explored for biological control purposes (e.g., Grundy and Maelzer 2000; Ambrose 2002). Among the exciting phenomena displayed by Harpactorinae are maternal care (Tallamy et al. 2004), specialization on termites in at least 2 lineages (Hwang and Weirauch 2012), the use of endogenous sticky secretions to aid in prey capture (Barth 1953; Zhang and Weirauch 2013b), and the collecting of plant resins to enhance prey capture and maternal care (Eisner 1988; Forero et al. 2011).



References

- Ambrose, D.P. 2002.** Assassin bugs (Heteroptera: Reduviidae) in integrated pest management programmes: success and strategies. *In* Strategies in Integrated Pest Management. *Edited by* S. Ignacimuthu and A. Sen. Phoenix Publishing House Pvt. Ltd., New Delhi, India. Pp. 73-85.
- Amyot, C.J.B. and Serville, A. 1843.** Histoire Naturelle des Insectes. Hemipteres. Roret, Paris.
- Barth, R. 1953.** Die Fangdrüsen an den Beinen von *Zelus (Diplocodus) leucogrammus* (Perty 1834) (Heteroptera, Reduviidae). *Zoologische Jahrbücher (Anatomie)*, **73**: 325-336.

Harpactorinae (cont.)

[Previous](#)

[Next](#)

References

- Bérenger, J. and Pluot-Sigwalt, D. 1997.** Special relationships of certain predatory Heteroptera Reduviidae with plants. First known case of a phytophagous Harpactorinae. *Comptes Rendus de l'Academie des Sciences Series III Sciences de la Vie*, **320**: 1007-1012.
- Davis, N.T. 1969.** Contribution to the morphology and phylogeny of the Reduvidae. Part IV. The harpactoroid complex. *Annals of the Entomological Society of America*, **62**: 74-94.
- Eisner T. 1988.** Insekten als fürsorgliche Eltern. *Verhandlungen der Deutschen Zoologischen Gesellschaft*, **81**: 9-17.
- Forero, D., Choe, D.H., and Weirauch, C. 2011.** Resin gathering in Neotropical resin bugs (Insecta: Hemiptera: Reduviidae): functional and comparative morphology. *Journal of Morphology*, **272**: 204-229.
- Forero, D., Berniker, L., and Weirauch, C. 2013.** Phylogeny and character evolution in the bee-assassins (Insecta: Heteroptera: Reduviidae). *Molecular Phylogenetics and Evolution*, **66**: 283-302.

Diaspidiini

Diaspidius scapha,
Gabon

© Jean-Michel Bérenger

Rhaphidosoma sp., Burkina Faso

© Jean-Michel Bérenger

Rhaphidosomini

Harpactorinae (cont.)

[Previous](#)

References (cont.)

- Grundy, P. and Maelzer, D. 2000.** Assessment of *Pristhesancus plagipennis* (Walker) (Hemiptera: Reduviidae) as an augmented biological control in cotton and soybean crops. *Australian Journal of Entomology*, **39**: 305-309.
- Miller, N.C.E. 1954.** The genus *Henricohahnia* Breddin, *Dicrotelus* Erichson, *Nyllius* Stål, *Orgetorixa* China and allied new genera. *Bulletin of the British Museum (Natural History). Entomology*, **3**: 447-485.
- Tallamy, D.W., Walsh, E., and Peck, D.C. 2004.** Revisiting paternal care in the assassin bug, *Atopozelus pallens* (Heteroptera: Reduviidae). *Journal of Insect Behavior*, **17**: 431-436.
- Tomokuni, M. and Cai, W. 2002.** *Asiacoris pudicus*, gen. & sp. n. of Dicrotelini (Heteroptera, Reduviidae, Harpactorinae) from Vietnam. *Deutsche Entomologische Zeitschrift*, **49**: 209-212.
- Zhang, G. and Weirauch, C. 2013a.** Molecular phylogeny of Harpactorini (Insecta: Reduviidae): correlation of novel predation strategy with accelerated evolution of predatory leg morphology. *Cladistics*, 1-13. doi: 10.1111/cla.12049.
- Zhang, G. and Weirauch, C. 2013b.** Sticky predators: a comparative study of sticky glands in harpactorine assassin bugs (Insecta: Hemiptera: Reduviidae). *Acta Zoologica*, **94**: 1-10.

Agriocoris flavipes,
Ecuador

Apiomerini

© Stephen Marshall

Calliclopius nigripes,
French Guiana

Apiomerini

© Jean-Michel Bérenger

Tegeini

Tegeini sp.,
Malaysia

© Melvyn Yeo

Holoptilinae

Diagnosis: Holoptilinae are sometimes referred to as feather-legged assassin bugs. They are diagnosed by the absence of the foretibial comb, presence of dense and long setae on the body and legs (Dasycnemini and Holoptilini), the trichome on the abdominal sternum (Holoptilini), and one closed cell (Holoptilini) or 2 longitudinal veins (Dasycnemini) on the forewing membrane. Feather-legged bugs in the tribe Aradellini share the general habitus of other Holoptilinae, but have short instead of long setation on legs and antenna.

Taxonomic Notes and Distribution: Holoptilinae comprise ~80 species in 16 extant and fossil (Poinar 1991) genera and 3 tribes (Holoptilini, Dasycnemini, and Aradellini; Wygodzinsky and Usinger 1963). The majority of species occur in the southern Palearctic, the Old World tropics and Australia (Malipatil 1985). One extant genus, *Neolocoptiris* Wygodzinsky and Usinger, is found in Guyana (Wygodzinsky and Usinger 1963).

Occurrence in Canada: Holoptilinae do not occur in Canada.

Natural History: Holoptilini appear to be specialized on ants and attract their prey with a seemingly glandular structure on the abdomen called the trichome (Weirauch and Cassis 2006; Weirauch et al. 2010).

[Back to Content](#)

[Back to Key](#)

Aradellus sp., Australia



Putoniola sp., Sudan



Smiliopus sp., Australia



Holoptilus ursus,
South Africa

© Christiane Weirauch

[Next](#)

Holoptilinae (cont.)

[Previous](#)

References

- Malipatil, M.B. 1985.** Revision of Australian Holoptilinae (Reduviidae: Heteroptera). *Australian Journal of Zoology*, **33**: 283-299.
- Poinar, G.O. 1991.** *Praecoris dominicana* gen. n., sp. n. (Hemiptera: Reduviidae: Holoptilinae) from Dominican amber, with an interpretation of past behavior based on functional morphology. *Insect Systematics and Evolution*, **22**: 193-199.
- Weirauch, C. and Cassis, G. 2006.** Attracting ants: the trichome and novel glandular areas on the sternum of *Ptilocnemus lemur* (Heteroptera: Reduviidae: Holoptilinae). *Journal of the New York Entomological Society*, **114**: 28-37.
- Weirauch, C., Bulbert, M., and Cassis, G. 2010.** Comparative trichome morphology in feather-legged assassin bugs (Insecta: Heteroptera: Reduviidae: Holoptilinae). *Zoologischer Anzeiger - A Journal of Comparative Zoology*, **248**: 237-253.
- Wygodzinsky, P.W. and Usinger, R.L. 1963.** Classification of the Holoptilinae and description of the first representative from the New World (Hemiptera: Reduviidae). *Proceedings of the Royal Entomological Society of London series B*, **32**: 47-52.



Ptilocnemus sp., Australia

Manangocorinae

[Back to Content](#)

[Back to Key](#)

Diagnosis: Manangocorinae are characterized by 2-segmented tarsi, the short and globose head that barely projects anteriorly beyond the small compound eyes, and the presence of a spine on the scutellum.

Taxonomic Notes and Distribution: This subfamily contains only one species described from Malaysia (Miller 1954). Whether this taxon merits subfamily status remains questionable (Schuh and Slater 1995). This is the most rarely collected subfamily of Reduviidae and is currently only known from the poorly preserved holotype.

Occurrence in Canada: Manangocorinae do not occur in Canada.


Natural History: The biology of Manangocorinae is unknown.

Reference

Miller, N.C.E. 1954. A new subfamily, new genera and species of Malaysian Reduviidae (Hem., Heteroptera). *Idea*, **10**: 1-8.

*Manangocoris
horridus*,
Malaysia

Holotype and
only known
specimen!



Peiratinae

Diagnosis: Peiratinae are characterized by the transverse sulcus on the pronotum located behind the middle of the pronotum, the typically rather elongate forecoxa with a distinctly flattened anterior surface, a typically very prominent fossula spongiosa on fore- and midtibiae, and the at least slightly asymmetrical male genitalia.

Taxonomic Notes and Distribution: The subfamily comprises ~350 species in ~34 genera. A number of peiratine genera were revised in modern times (e.g., Coscarón 1983, 1997; Willemse 1985; Zhang and Weirauch 2011). Peiratinae were supported as one of the early lineages within the Higher Reduviidae and are thus likely a rather old (~100 million years) group of Reduviidae (Hwang and Weirauch 2012). Peiratinae have cosmopolitan distribution.

Occurrence in Canada: One species, *Melanolestes picipes* (Herrich-Schäffer), is known to occur in Canada.

Natural History: Members of this subfamily are primarily ground-dwelling or found on decomposing tree trunks and in other cryptic microhabitats. They are usually nocturnal and thus frequently attracted to lights. Peiratinae are also known for their extremely painful bite (Radio 1927). Male Peiratinae are unique among Reduviidae in possessing slightly to strongly asymmetrical genitalic structures (e.g., Willemse 1985). Males in some species also feature “extragenital structures” that may be involved in mating (Ghuri 1964). Some peiratine species mate end-to-end (Haridass 1985; Ambrose 1999), a mating position that is otherwise unknown in Reduviidae.

[Back to Content](#)

[Back to Key](#)



[Next](#)

Peiratinae (cont.)

[Previous](#)

References

Ambrose, D.P. 1999. Assassin Bugs. Science Publishers Inc., Enfield.

Coscarón, M. C. 1983. Revision del genero *Rasahus* (Insecta, Heteroptera, Reduviidae). Revista del Museo de La Plata Zoologia, **13**: 75-138.

Coscarón, M.C. 1997. Revision of the genus *Peirates* Serville, with a cladistic and biogeographic analysis (Heteroptera: Reduviidae, Peiratinae). Insect Systematics and Evolution, **28**: 39-73.

Ghuri, M.S.K. 1964. A remarkable phenomenon amongst the males of Piratinae (Reduviidae, Heteroptera). The Annals and Magazine of Natural History, **7**: 733-737.

Haridass, E.T. 1985. Feeding and ovipositional behavior in some reduviids (Insecta-Heteroptera). Proceedings of the Indian Academy of Sciences, **94**: 239-247.

Radio, P.A. 1927. Studies on the biology of the Reduviidae of American north of Mexico. The University of Kansas Science Bulletin, **17**: 5-291.

Willemse, L. 1985. A taxonomic revision of the New World species of *Sirthenea* (Heteroptera: Reduviidae: Peiratinae). Zoologische Verhandelingen, **215**: 1-67.

Zhang, G. and Weirauch, C. 2011. Matching dimorphic sexes and immature stages with adults: resolving the systematics of the *Bekilya* group of Malagasy assassin bugs (Hemiptera: Reduviidae: Peiratinae). Systematic Entomology, **36**: 115-138.



Phimophorinae

Diagnosis: The Phimophorinae are characterized by an aradid-like body that is covered with waxy secretions, the head almost twice as long as high, incrassate antennal segments, buccula obscuring the base of the labium, the second (first visible) labial segment much longer than segments 3 and 4 combined, the largely membranous forewings, the small 2-segmented tarsi, and the prosternum with shield-like structures near the stridulatory groove.

Taxonomic Notes and Distribution: This cryptic and rarely collected subfamily currently only contains the genera *Mendanocoris* Miller (2 species; Solomon Islands and Malaysia) and *Phimophorus* Bergroth (one species; Neotropics) [Usinger and Wygodzinsky 1964; Schuh and Slater 1995]. The phylogenetic position of this group has been the focus of much debate throughout the 20th century (Handlirsch 1897a, 1897b; Usinger and Wygodzinsky 1964). Maldonado (1990) recognized 2 tribes, the Mendanocorini Miller and Phimophorini Handlirsch, a scheme we do not follow in the current study. Due to the great rarity of specimens in collections, Phimophorinae have not been included in recently published phylogenies of Reduviidae, making the analysis of the phylogenetic position of this group one of the top priorities in reduviid systematics.

Occurrence in Canada: Phimophorinae do not occur in Canada.

[Back to Content](#)

[Back to Key](#)



Phimophorus spissicornis, Colombia

Phimophorinae (cont.)

[Previous](#)

Natural History: The natural history of Phimophorinae was virtually unknown until recently. Fieldwork in Colombia revealed that *Phimophorus spissicornis* Bergroth is typically associated with palms (family Arecaceae), but may also occur in bromeliads and bananas (Chaverra-Rodriguez et al. 2010). Specimens were collected in association with *Rhodnius pallescens* Barber, confirming earlier observations that *Phimophorus* may co-occur with Triatominae (Lent and Jurberg 1977; Lent and Wygodzinsky 1979). An association with palm trees and specimens of *Rhodnius prolixus* was also observed in Ecuador (Bérenger, unpublished data).

References

- Handlirsch, A. 1897a.** Ueber *Phimophorus spissicornis* Bergr. Ein hemipterologischer Beitrag. Verhandelingen Zoologischbotanischen Gesellschaft in Wien, **47**: 408-410.
- Handlirsch, A. 1897b.** Zur systematischen Stellung der Hemipterengattung *Phimophorus* Bergroth. Verhandelingen Zoologischbotanischen Gesellschaft in Wien, **47**: 558-560.
- Chaverra-Rodriguez, D., Forero, D., Jaramillo O.N., and Triana Ch, O. 2010.** New record and ecological notes of *Phimophorus spissicornis* (Hemiptera: Heteroptera: Reduviidae: Phimophorinae) in Colombia. Revista Colombiana de Entomologia, **36**: 176-178.
- Lent, H. and Jurberg, J. 1977.** As ninfas de quinto estágio de *Phimophorus spissicornis* Bergroth, 1886 (Hemiptera, Reduviidae). Revista Brasileira de Biología, **37**: 521-524.
- Lent, H. and Wygodzinsky, P.W. 1979.** Revision of the Triatominae (Hemiptera, Reduviidae), and their significance as vectors of Chagas disease. Bulletin of the American Museum of Natural History, **163**: 123-520.
- Usinger, R.L. and Wygodzinsky, P. 1964.** Description of a new species of *Mandanocoris* Miller, with notes on the systematic position of the genus (Reduviidae, Hemiptera, Insect). American Museum Novitates, **2204**: 1-13.

Phymatinae

[Back to Content](#)

[Back to Key](#)

Diagnosis: Commonly called ambush bugs, Phymatinae are a group of morphologically very distinctive assassin bugs. Most members of the subfamily are characterized by the presence of buccula obscuring the base of the labium and the head often being about as high as long. The conspicuously enlarged forefemur is diagnostic for Macrocephalini and Phymatini and the chelate or subchelate foreleg is characteristic for Carcinocorini, Macrocephalini, and Phymatini. The foreleg in Themonocorini is shaped like a standard walking leg with the femur featuring long and stout spines.

Taxonomic Notes and Distribution: Phymatinae were considered as a distinct family within Cimicomorpha by several authors (Froeschner and Kormilev 1989; Maldonado 1990), and the name Phymatidae is still being used in some current literature and popular websites. However, Carayon et al. (1958) made a very convincing case for treating Phymatinae as part of the Reduviidae, based on shared unique features with several other reduviid subfamilies. All recent cladistic analyses have confirmed Phymatinae to be part of the “phymatine complex” of reduviid subfamilies that also contains the Centrocnemidinae, Elasmodeminae, and Holoptilinae (Weirauch 2008; Weirauch and Munro 2009; Hwang and Weirauch 2012). Monographs on Phymatinae include Handlirsch (1897), Kormilev (1962), and Maa and Lin (1959). Four tribes are currently recognized: Carcinocorini (Oriental), Macrocephalini (cosmopolitan), Phymatini (cosmopolitan), and Themonocorini (Afrotropical). Froeschner and Kormilev (1989) provided a catalogue of Phymatinae, recording 281 species and 26 genera. Since then, 2 new genera, 10 new species, and one new subspecies have been described (Kormilev 1990a, 1990b, 1990c; Kormilev and van Doesburg 1991, 1992; van Doesburg 1997, 2004; Cui et al. 2003; Rabitsch 2003; Rabitsch et al. 2006; van Doesburg and Pluot-Sigwalt 2007; van Doesburg and Jacobs 2011).

Occurrence in Canada: Some species in the genus *Phymata* Latreille are known to occur in Canada.

[Next](#)

Phymatinae (cont.)

[Previous](#)

[Next](#)



Themonocoris endroedyi, South Africa



© Christiane Weirauch

Macrocephalus sp., United States

Natural History: Phymatinae typically hide among flowers or vegetation, and mount an ambush attack on their prey that consists of a variety of small arthropods. Adults and nymphs of the African *Themonocoris kinkalanus* Carayon, Usinger, and Wygodzinsky were observed to seek their prey in nests of the weaver bird genus *Ploceus* Cuvier, in flower and fruit clusters of the palm genus *Elaeis* Jacquin, and in dead plant matter (Carayon et al. 1958). Pre- and postcopulatory mate guarding behaviors occur in *Phymata fasciata* (Gray) [Dodson and Marshall 1984], production of acoustical signals has been recorded for *Phymata crassipes* (Fabricius) [Gogala and Cokl 1983], and species of *Phymata* have recently become the focus of sexual selection studies (e.g., Punzalan et al. 2008a, 2008b).

Phymata spp., various countries



© Amy Michael and Sarah Frankenberg

Phymatinae (cont.)

[Previous](#)[Next](#)

References

- Carayon, J., Usinger, R.L., and Wygodzinsky, P.W. 1958.** Notes on the higher classification of the Reduviidae, with the description of a new tribe of the Phymatinae (Hemiptera-Heteroptera). *Revue de Zoology et Botanique Africaine*, **57**: 256-281.
- Cui, J., Cai, W., and Rabitsch, W. 2003.** Notes on the genus *Agdistocoris* Kormilev, 1962 (Heteroptera: Reduviidae: Phymatinae), with the description of a new species from China. *Annales Zoologici (Warszawa)*, **53**: 669-671.
- Dodson, G. and Marshall, L. 1984.** Mating Patterns in an Ambush Bug *Phymata fasciata* (Phymatidae). *The American Midland Naturalist*, **112**: 50-57.
- Froeschner, R.C. and Kormilev, N.A. 1989.** Phymatidae or ambush bugs of the world: a synonymic list with keys to species, except *Lophoscutus* and *Phymata* (Hemiptera). *Entomography*, **6**: 1-76.
- Gogala, M. and Cokl, A. 1983.** The acoustical behavior of the bug *Phymata crassipes* (F.) (Heteroptera). *Revue Canadienne de Biologie Experimentale*, **42**: 249-256.
- Handlirsch, A. 1897.** Monographic der Phymatiden. *Annalen des Kaiserlich-königlichen Naturhistorischen Hofmuseum*, **12**: 127-230.
- Kormilev, N.A. 1962.** Revision of Phymatinae (Hemiptera, Phymatidae). *Philippine Journal of Science Manila*, **89**: 287-486.
- Kormilev, N.A. 1990a.** Notes on American Phymatidae, with description of a new species of *Phymata* Latreille. *Anales del Instituto de Biologia Universidad Nacional Autonoma de Mexico Serie Zoologia*, **61**: 99-106.
- Kormilev, N.A. 1990b.** Two new species of the genus *Lophoscutus* Kormilev, 1951 (Hemiptera, Phymatidae). *Revista Brasileira de Entomologia*, **34**: 221-225.
- Kormilev, N.A. 1990c.** A new species of the genus *Chelocoris* from Vietnam (Hemiptera: Phymatidae: Carcinocorinae). *Bishop Museum Occasional Papers*, **30**: 298-300.

Phymatinae (cont.)

[Previous](#)[Next](#)

References (cont.)

- Kormilev, N.A. and van Doesburg, P.H. 1991.** Notes on Caribbean Phymatidae (Hemiptera: Heteroptera). *Zoologische Mededeelingen*, **65**: 277-285.
- Kormilev, N.A. and van Doesburg, P.H. 1992.** Notes on Macrocephalinae (Heteroptera: Phymatidae). *Zoologische Mededeelingen*, **66**: 417-421.
- Maa T.C. and Lin, K.S. 1959.** Further notes on Old World Phymatinae (Hemiptera: Reduviidae). *Pacific Insects*, **1**: 315-323.
- Punzalan, D., Rodd, F.H., and Rowe, L. 2008a.** Contemporary sexual selection on sexually dimorphic traits in the ambush bug *Phymata americana*. *Behavioral Ecology*, **19**: 860-870.
- Punzalan, D., Rodd, F.H., and Rowe, L. 2008b.** Sexual selection mediated by the thermoregulatory effects of male colour pattern in the ambush bug *Phymata americana*. *Proceedings of the Royal Entomological Society London Series B*, **275**: 483-492.
- Rabitsch, W. 2003.** *Glossopelta laotica* sp. n. (Insecta: Heteroptera: Reduviidae: Phymatinae), a new ambush bug from Laos. *Annalen des Naturhistorischen Museums in Wien Serie B Botanik und Zoologie*, **105B**: 447-451.
- Rabitsch, W., Cui, J., and Cai, W. 2006.** Two new ambush bugs from Yunnan, southern China (Insecta, Heteroptera, Reduviidae, Phymatinae). *Denisia*, **19**: 581-588.
- van Doesburg, P.H. 1997.** A case of homonymy in the Phymatidae (Heteroptera). *Entomologische Berichten (Amsterdam)*, **57**: 96.
- van Doesburg, P.H. 2004.** *Goellneriana deckerti* gen. nov. & spec. nov. (Heteroptera: Reduviidae: Phymatinae: Macrocephalini) from Namibia, with a review of the Afrotropical ambush bugs. *Zoologische mededeelingen*, **78**: 147-159.

Phymatinae (cont.)

[Previous](#)

References (cont.)

van Doesburg, P.H. and Jacobs, D.H. 2011. A new *Themonocoris* from South Africa (Hemiptera: Heteroptera: Reduviidae: Phymatinae: Themonocorini). *Zoologische Mededeelingen*, **85**: 55-67.

van Doesburg, P.H. and Pluot-Sigwalt, D. 2007. Notes on Phymatinae (Insecta: Heteroptera: Reduviidae). The genus *Oxythyreus* Westwood, 1841. *Zoologische Mededeelingen*, **81**: 85-99.

Weirauch, C. 2008. Cladistic analysis of Reduviidae (Heteroptera: Cimicomorpha) based on morphological characters. *Systematic Entomology*, **33**: 229-274.

Weirauch, C. and Munro, J.B. 2009. Molecular phylogeny of the assassin bugs (Hemiptera: Reduviidae), based on mitochondrial and nuclear ribosomal genes. *Molecular Phylogenetics and Evolution*, **53**: 287-299.

Glossopelta sp., China



Phymata cf. *americana*,
United States



Phymata
americana,
Canada



Physoderinae

Diagnosis: Species of Physoderinae are characterized by a tuberculate body with spatulate setae, the foretibial comb located on a subapical spur, a usually elongate head with small compound eyes that are far from the anterior margin of the pronotum, the third labial segment (second visible) very long and straight, the forewing membrane with 2 cells, the rounded forecoxa, and the absence of a fossula spongiosa.

Taxonomic Notes and Distribution: The subfamily status of Physoderinae was established by Miller (1954). The group currently comprises >60 species in ~15 genera. Villiers (1962) monographed the fauna of Madagascar, one of the hotspots of physoderine biodiversity. *Cryptophysoderes fairchildi* Wygodzinsky and Maldonado was described as the first Neotropical representative of this otherwise Old World clade (Wygodzinsky and Maldonado 1972), followed by the description of a second monotypic New World genus, *Leptophysoderes* Weirauch (Weirauch 2006). Despite its diversity in Madagascar, Physoderinae were not known to occur on the African mainland until the discovery of a cryptic, apterous taxon from Tanzania that was described as *Porcelloderes* Rédei (Rédei 2012).

Occurrence in Canada: Physoderinae do not occur in Canada.

Natural History: Specimens of the subfamily have been found in caves, hollow trees, vegetable debris, and at the bases of banana and *Pandanus* plants (Schuh and Slater 1995).

[Back to Content](#)

[Back to Key](#)

Physoderes notata,
Indonesia



Breviphysoderes decora,
Brunei



© Christiane Weirauch

[Next](#)

Physoderinae

[Previous](#)

References

- Miller, N.C.E. 1954.** New genera and species of Reduviidae from Indonesia and the description of a new subfamily (Hemiptera-Heteroptera). *Tijdschrift Voor Entomologie*, **97**: 76-114.
- Rédei, D. 2012.** *Porcelloderes impenetrabilis* gen. & sp. n. from Tanzania, an Assassin Bug Camouflaging in the Adult Stage (Hemiptera: Heteroptera: Reduviidae: Physoderinae). *African Invertebrates*, **53**: 601-613.
- Villiers, A. 1962.** Les Reduviides de Madagascar XX, Physoderinae. *Revue Francaise d'Entomologie*, **29**: 219-234.
- Weirauch, C. 2006.** New Genus and Species of Physoderinae (Heteroptera: Reduviidae) from the New World, with a Revised Diagnosis of Physoderinae Miller. *American Museum Novitates*, **3510**: 1-9.
- Wygodzinsky, P.W. and Maldonado, J.C. 1972.** Description of the first genus of physoderine assassin bugs (Reduviidae, Hemiptera) from the New World. *American Museum Novitates*, **2504**: 1-7.



Pseudocetherinae

Diagnosis: Pseudocetherinae are sometimes referred to as lobe-headed bugs. Members of this subfamily are characterized by the short mandibular plates with the apices well separated by the elongate clypeus; the maxillary plates with a large anterior projection giving the head a tri-lobed appearance; the forewing membrane with a small triangular Cu-PCu cell; the first visible labial segment long and slender, longer than the second and third segments combined; the antennal scape short and not or only slightly surpassing the apex of the head; the prosternum projecting anteriorly, forming a nearly ventrally flattened surface; and the dorsal abdominal glands entirely absent in adults.

Taxonomic History and Distribution: In 1963, Villiers described the genus *Pseudocethera* and placed it in a new subfamily, the Pseudocetherinae. The systematic position of the genus, which currently contains 3 Afrotropical species, was modified subsequently by different authors, some treating it as part of the Cetherinae (Putchkov and Putchkov 1985), others retaining it as a separate subfamily. Species of *Pseudocethera* were not included in recent phylogenetic analyses of Reduviidae. However, several genera currently treated as part of the (polyphyletic) subfamily Reduviinae share many of the diagnostic features listed above, among them *Kayanocoris* Miller and part of the genus *Gerbelius* Distant (Weirauch and Rédei, unpublished). Hwang and Weirauch (2012) found these 2 genera to be part of the *Reduvius* clade of Reduviinae, refuting a close relationship with the Cetherinae. We here treat Pseudocetherinae as a distinct subfamily, pending the inclusion of several additional genera.



Pseudocethera sp., Kenya

© Christiane Weirauch

Pseudocetherinae (cont.)

[Previous](#)

Occurrence in Canada: Pseudocetherinae do not occur in Canada.

Natural History: The biology of *Pseudocethera* is unknown.

Reference

Villiers, A. 1963. Hemiptera Reduviidae. La reserve naturelle integrale du Mont Nimba. XXV Hemiptere Reduviidae. *Memoirs de l'Instiut francais d'Afrique Noir*, **66**: 479-565.

Pseudocethera sp., Kenya



© Christiane Weirauch



Reduviinae

[Back to Content](#) [Back to Key](#)

Diagnosis: Reduviinae are recognized by usually having ocelli, 3-segmented tarsi, 2 closed cells on the forewing membrane, and a fossula spongiosa on fore- and midtibiae. In addition, Reduviinae are typically recognized by the absence of specialized morphological features that are characteristic for other subfamilies.

Taxonomic History and Distribution: At least 140 genera and ~1,000 species are currently classified as Reduviinae. Species of Reduviinae are primarily recognized by the absence of characters found in other reduviids (Schuh and Slater 1995), a fact that is not surprising given the polyphyletic nature of this assemblage of unrelated groups (Hwang and Weirauch 2012). The extent of the “reduviine polyphyly problem” is currently being investigated, and the most recent, molecular-only phylogenetic analysis of Reduviidae (Hwang and Weirauch 2012) indicates that Reduviinae may fall into at least 11-13 distinct monophyletic clades. Reduviinae, as currently classified, are cosmopolitan, with the greatest diversity in the Old and New World tropics.

Occurrence in Canada: Only the cosmopolitan species *Reduvius personatus* (Linnaeus) occurs in Canada.



Reduviinae (cont.)

[Previous](#)

[Next](#)

Natural History: Most species are nocturnal and believed to be generalist predators of insects and other arthropods (Schuh and Slater 1995). Many Reduviinae are found in association with the bark of either living or dead trees (Miller 1953), some species are specialized ant predators (*Acanthaspis* spp.; Louis 1974; Haridass 1985), and some species may live in animal burrows and nests (e.g., *Reduvius* spp. and *Opisthacidius* spp.; Lent and Wygodzinsky 1956; Ryckmann 1954). The cosmopolitan *R. personatus* is closely associated with humans: adults are attracted to lights and camouflaged immatures can be found in houses.



Reduviinae (cont.)

[Previous](#)

References


Haridass, E.T. 1985. Feeding and ovipositional behavior in some reduviids (Insecta-Heteroptera). Proceedings of the Indian Academy of Sciences, **94**: 239-247.

Lent, H. and Wygodzinsky, P.W. 1956. Situação atual do gênero *Opisthacidius*. Revista Brasileira de Biologia, **16**: 327-334.

Louis, D. 1974. Biology of Reduviidae of cocoa farms in Ghana. American Midland Naturalist, **91**: 68-89.

Miller, N.C.E. 1953. Notes on the biology of the Reduviidae of southern Rhodesia. Transactions of the Zoological Society of London, **27**: 541-672.

Ryckman, R.E. 1954. *Reduvius senilis* Van Duzee from the lodges of *Neotoma* in San Juan county, Utah (Hemiptera: Reduviidae). Bulletin of the Southern California Academy of Sciences, **53**: 88.



Saicinae

[Back to Content](#) [Back to Key](#)

Diagnosis: Saicinae are diagnosed by the absence of ocelli and fossula spongiosa, the second visible labial segment frequently expanded and basally bulbous, the forecoxa at most 3 times as long as wide, the usually enlarged and slightly bulbous tarsal segments, the acetabulum of the foreleg oriented ventrad, and the opposing surfaces of head and labium frequently with stiff setae or spines.

Taxonomic History and Distribution: Saicinae were first recognized by Stål (1859). The subfamily currently comprises ~25 genera and >140 species. Wygodzinsky (1966) treated Saicinae as closely related to the Emesinae and Visayanocorinae, and recent phylogenetic analyses have confirmed this relationship (e.g., Hwang and Weirauch 2012). Gil-Santana and Costa (2009) and Villiers (1969) produced keys to the American and African genera, respectively.

Occurrence in Canada: Saicinae have not been recorded from Canada.

Natural History: Specimens of this subfamily are commonly collected at lights. Little else is currently known of Saicinae life history.

Saica apicalis, French Guiana



© Jean-Michel Bérenger

[Next](#)

Saicinae (cont.)

[Previous](#)

References

- Gil-Santana, H.R. and Costa, L.A.A. 2009.** A new species of *Paratagalis* Monte from Brazil with taxonomical notes and a key to New World genera of Saicinae (Hemiptera: Heteroptera: Reduviidae: Saicinae). *Zootaxa*, **2197**: 20-30
- Stål, C. 1859.** Nova methodus Reduvina (Burm.) disponendi. *Berliner Entomologische Zeitschrift*, **3**: 328.
- Villiers, A. 1969.** Revision des Reduviides africains. IV. Saicinae. *Bulleting de l'Institut Fondamental d'Afrique Noire*, **31**: 1186-1247.
- Wygodzinsky, P.W. 1966.** A monograph of the Emesinae (Reduviidae, Hemiptera). *Bulletin of the American Museum of Natural History*, **133**: 1-614.



Salyavatinae

[Back to Content](#) [Back to Key](#)

Diagnosis: Salyavatinae are characterized by a 2-segmented foretarsus, fossula spongiosa on the fore- and midtibiae, and a small, short head with antennal tubercles at the base of the antennae. Most members possess various spines on the head, pronotum, scutellum, postscutellum (metanotum), legs, and abdomen. Almost all species of this group are cryptically colored (primarily brown, some with pale and dark marmorated patterns). Members of the largest genus *Lisarda* Stål have a frontal process, while members of several African and Oriental genera have variously inflated foretibiae.

Taxonomic History and Distribution: Salyavatinae were classified as a subfamily by Amyot and Serville (1843). Currently, the subfamily consists of 17 genera and 108 species but the group is in need of revision. Van Doesburg and Forero (2012) revised the genus *Salyavata* Amyot and Serville. With the exception of the Neotropical genus *Salyavata*, all species occur in the Old World tropics.

Occurrence in Canada: Salyavatinae do not occur in Canada.

Natural History: Many genera are known to camouflage as nymphs and all members of this group are suspected to feed on termites (Miller 1953), with the genus *Salyavata* being the best studied (McMahan 1982, 1983a, 1983b). The function of the inflated foretibial structures is unknown.

Lisarda pallidispina, Malaysia



© Melvyn Yeo

Lisarda inornata, Singapore



© Melvyn Yeo

[Next](#)

Salyavatinae (cont.)

[Previous](#)

References

Amyot, C.J.B. and Serville, A. 1843. Histoire Naturelle des Insectes. Hemipteres. Roret, Paris.

McMahan, E.A. 1982. Bait-and-capture strategy of a termite-eating assassin bug. *Insectes Sociaux*, **29**: 346–351.

McMahan, E.A. 1983a. Adaptations, feeding preferences, and biometrics of a termite-baiting assassin bug (Hemiptera: Reduviidae). *Annals of the Entomological Society of America*, **76**: 483-486.

McMahan, E.A. 1983b. Bugs angle for termites. *Natural History*, **92**: 40-47.

Miller, N.C.E. 1953. Notes on the biology of the Reduviidae of southern Rhodesia. *Transactions of the Zoological Society of London*, **27**: 541-672.

van Doesburg P.H. and Forero, D. 2012. Revision of the genus *Salyavata* Amyot & Audinet-Serville, 1843 (Heteroptera: Reduviidae: Salyavatinae). *Zoologische Mededeelingen*, **86**: 515-568.

Salyavata variegata,
Ecuador



© Stephen Marshall

Valentia compressipes, Vietnam



© Stephen Marshall

Sphaeridopinae

Diagnosis: Sphaeridopinae are characterized by a large, robust body; large eyes almost covering the entire head; and a short, straight, thin labium. They share the presence of antennal tubercles with the Salyavatinae but differ in possessing a 3-segmented foretarsus. *Sphaeridops* Amyot and Serville and *Volesus* Champion are mostly red and black, whereas species of *Veseris* Stål are brown marmorated.

Taxonomic History and Distribution:

Sphaeridopinae were classified as a family by Pinto (1927), and as a subfamily by Costa Lima (1940), and currently consist of 3 genera and 6 species (Maldonado and Santiago-Blay 1992; Gil-Santana et al. 2000; Gil-Santana and Alencar 2001). They are only found in the Neotropics.

Occurrence in Canada: Sphaeridopinae do not occur in Canada.

Natural History: Nymphs of Sphaeridopinae have camouflaging structures, have been found in association with termites, and have been fed on termites in captivity (Wygodzinsky, pers. com. in: McMahan 1982).

[Back to Content](#)

[Back to Key \(taxa without fossula\)](#)

[Back to Key \(taxa with fossula\)](#)



Veseris bellator, French Guiana

© Jean-Michel Bérenger

[Next](#)

Sphaeridopinae (cont.)

References

- Costa Lima, A. 1940.** Insetos do Brasil , 2. Hemipteros. Escola Nacional de Agronomia, ser. Didáctica nº3, 2, 351 pp
- Gil-Santana, H.R., Costa, L.A.A., and Zeraik, S.O. 2000.** A new species of *Sphaeridops* Amyot & Serville, 1843 (Hemiptera, Reduviidae, Sphaeridopinae). [Especie nova de *Sphaeridops* Amyot & Serville, 1843 (Hemiptera, Reduviidae, Sphaeridopinae)] Boletim do Museu Nacional de Zoologia, **423**: 1-6
- Gil-Santana, H.R. and Alencar, J. 2001.** Sobre o gênero *Veseris* Stål, 1865, com *Eurylochus* Torre- Bueno, 1914, como sinônimo novo e chaves para identificação (Hemiptera, Reduviidae, Sphaeridopinae). Entomología y Vectores, **8**: 95-104.
- Maldonado, C.J. and Santiago-Blay, J.A. 1992.** A new species of the Neotropical genus *Sphaeridops* Amyot & Serville, 1843 (Sphaeridopinae: Reduviidae). Proceedings of the Entomological Society of Washington, **94**: 508-511.
- McMahan, E.A. 1982.** Bait-and-capture strategy of a termite-eating assassin bug. Insectes Sociaux, **29**: 346-351.
- Pinto, C. 1927.** Sphaeridopidae, nova familia de Hemiptero Reduvidae, com a descrição de um gênero e especie nova. Boletim Biologico, Sao Paulo, **6**: 43-51.

[Previous](#)



Volesus nigripennis,
Colombia

© Dimitri Forero



Sphaeridops amoenus,
Brazil

© Christiane Weirauch

Stenopodainae


[Back to Content](#) [Back to Key](#)

Diagnosis: Stenopodainae are a cryptically colored group of assassin bugs characterized by a pentagonal or hexagonal cell in the corium. The mandibular plates are prominent, and the pedicel and flagellomeres of the antennae are often folded backwards under the scape.

Taxonomic History and Distribution: This is a large group of more than 110 genera (Maldonado 1990). The group is worldwide with most species occurring in the tropics. Barber (1929) published a first synopsis on the subfamily, Wygodzinsky and Giacchi (1991) provided a key for Neotropical genera and Giacchi (e.g., 1969, 1998) revised the Neotropical genera in a series of 12 papers. Stenopodainae of Madagascar were studied by Villiers (1968).

Occurrence in Canada: Stenopodaine genera recorded from Canada are *Phirontis* Stål, *Pygolampis* Germar, and *Stenopoda* Laporte.

Natural History: Stenopodainae are often found in leaf litter or soil, but many specimens have also been collected at lights. Numerous species are apterous. Stenopodainae are also commonly found on islands (Rédei 2005; Chlond 2010).



Oncocephalus sp., Namibia

© Stephen Marshall



Stenopodainae sp., Vietnam

© Stephen Marshall

[Next](#)

Stenopodainae (cont.):

[Previous](#)

References

- Barber H.G. 1929.** Essay on the subfamily Stenopodainae of the new world. *Entomologica Americana*, X, **3**: 149-238.
- Chlund, D. 2010.** *Planeocoris*, a new genus of Stenopodainae (Hemiptera: Heteroptera: Reduviidae) from Madagascar. *Zootaxa*, **2400**: 41-48.
- Giacchi, J.C. 1969.** Revision del genero *Stenopoda* Laporte, 1833 (Hemiptera, Reduviidae, Stenopodainae). *Physis*, **29**: 1-26.
- Giacchi, J.C. 1998.** Revision de los Stenopodainos americanos. XI. Relaciones entre los generos *Ctenotrachelus* Stal, 1868, *Seridentus* Osborn, 19040 y *Achillas* Torre Bueno, 1914 (Heteroptera, Reduviidae). *Physis*, **56**: 27-30.
- Paiero, S.M., Marshall, S.A., and Hamilton, K.G.A. 2004.** New records of Hemiptera from Canada and Ontario. *Proceedings of the Entomological Society of Ontario*, **134**: 115-129.
- Rédei, D. 2005.** New synonymy of the endemic reduviid on St. Helena Island (Heteroptera: Reduviidae, Stenopodainae). *Entomologist's Monthly Magazine*, **141**: 93-94.
- Wygodzinsky, P.W. and Giacchi, J. C. 1991.** Key to the genera of Stenopodainae of the new world (Insecta, Heteroptera, Reduviidae). *Physis*, **49**: 5-9.

Canthesancus gulo, Singapore



© Melvyn Yeo

Sastrapada sp.,
Singapore



© Melvyn Yeo

*Oncocephalus
cingalensis*,
Singapore



© Melvyn Yeo

Triatominae

[Back to Content](#)

[Back to Key \(taxa without fossula\)](#)

[Back to Key \(taxa with fossula\)](#)

Diagnosis: Triatominae, or kissing bugs, are characterized by a long, straight third (second visible) labial segment with flexible membranous connection between the last 2 segments. Most members have an elongate head with no constriction behind the eyes and the ocelli elevated on a protuberance. The connexivum sometimes has membranous areas to allow for expansion upon blood feeding.

Taxonomic History and Distribution: Lent and Wygodzinsky (1979) monographed Triatominae, and this publication is still an important reference. The subfamily consists of 5 tribes (Rhodniini, Cavernicolini, Bolboderini, Alberproseniini, Triatomini) containing 16 genera and ~140 species (Galvão et al. 2003). Most are found in the New World but one genus, *Linshcosteus* Distant, is restricted to India and several species of *Triatoma* Laporte are found throughout the Oriental region. Even though Triatominae have been treated as a distinct subfamily for a long time, recent phylogenetic research has cast doubt on the monophyly of this group. Published hypotheses currently find support for either the monophyly of Triatominae (Hypša et al. 2002; Weirauch and Munro 2009; Patterson and Gaunt 2010), their polyphyly (Paula et al. 2005; Schofield and Galvão 2009), or their paraphyly (Hwang and Weirauch, 2012) with respect to several genera of large-bodied Reduviinae.

Triatoma dimidiata, Costa Rica



Panstrongylus geniculatus,
French Guiana



Rhodnius robustus, French Guiana



[Next](#)

Triatominae (cont.)

[Previous](#)

Occurrence in Canada: Triatominae have not been recorded from Canada.

Natural History: Members of this group feed on vertebrate blood and many species are vectors of Chagas disease that is caused by the parasite *Trypanosoma cruzi* (Chagas) [Lent and Wygodzinsky 1979].

References

- Galvão, C., Carcavallo, R.U., Rocha, D.S., and Jurberg, J. 2003.** A checklist of the current valid species of the subfamily Triatominae Jeannel, 1919 (Hemiptera, Reduviidae) and their geographical distribution, with nomenclatural and taxonomic notes. *Zootaxa*, **202**: 1-36.
- Hypša, V., Tietz, D.F., Zrzavy, J., Rego, R.O.M, Galvão, C., and Jurberg, J. 2000.** Phylogeny and biogeography of Triatominae (Hemiptera: Reduviidae): molecular evidence of a New World origin of the Asiatic clade. *Molecular Phylogenetics and Evolution*, **23**: 447-457.
- Lent, H. and Wygodzinsky, P.W. 1979.** Revision of the Triatominae (Hemiptera, Reduviidae), and their significance as vectors of Chagas' disease. *Bulletin of the American Museum of Natural History*, **163**: 123-520.
- Patterson, J.S. and Gaunt, M.W. 2010.** Phylogenetic multi-locus codon models and molecular clocks reveal the monophyly of haematophagous reduviid bugs and their evolution at the formation of South America. *Molecular Phylogenetics and Evolution*, **56**: 608-621.
- Paula, A.S. de, Diotaiuti, L., and Schofield, C.J. 2005.** Testing the sister-group relationship of the Rhodniini and Triatomini (Insecta: Hemiptera: Reduviidae: Triatominae). *Molecular Phylogenetics and Evolution*, **35**: 712–718.
- Schofield, C.J. and Galvão, C. 2009.** Classification, evolution, and species groups within the Triatominae. *Acta Tropica*, **110**: 88-100.
- Weirauch, C. and Munro, J.B. 2009.** Molecular phylogeny of the assassin bugs (Hemiptera: Reduviidae), based on mitochondrial and nuclear ribosomal genes. *Molecular Phylogenetics and Evolution*, **53**: 287-299.

Tribelocephalinae

Diagnosis: Tribelocephalinae are characterized by dense specialized setae with a bulbous base and curved apex that cover much of the body. In almost all members, the ocelli are absent, the eyes are flat, and the scape is stout and longer than the head, with the more distal segments folded backwards underneath it. In addition, the corium is reduced such that the membrane takes up most of the hemelytron. Members are often brown, cryptic and slightly flattened.

Taxonomic History and Distribution: The group consists of 16 genera and ~130 species (Maldonado 1990; Rédei 2007; Weirauch 2010). Tribelocephalinae were first recognized by Stål (1866). Villiers (1943) divided the group into 2 tribes, the Opistoplatyini and Tribelocephalini. Maldonado (1996) created a third tribe, the Xenocaucini. The genus *Xenocaucus* China and Usinger, with 2 species, is known from Bioko Island in the Gulf of Guinea and from Tanzania. It is eyeless, wingless, and covered in dense setae (China and Usinger 1949). Tribelocephalinae occur in the Old World tropics with exception of the recently described genus *Tribelocodia* Weirauch that is found in French Guiana. *Tribelocodia* possesses some features otherwise typically found in Ectrichodiinae (ocelli, 2-pronged scutellum, and fossula spongiosa), suggesting that either the monophyly of these 2 closely related taxa is questionable, or that character evolution within these groups is more complex than previously assumed (Weirauch 2008; Weirauch 2010; Hwang and Weirauch 2012).

[Back to Content](#)

[Back to Key](#)

Opistoplatys sp.,
Indonesia



[Next](#)

© Christiane Weirauch

Abelocephala
sp., Australia



[Previous](#)

Tribelocephalinae (cont.)

Occurrence in Canada: Tribelocephalinae do not occur in Canada.

Natural History: Tribelocephalinae are often found in leaf litter and some are attracted to lights, but their biology is largely unknown (Schuh and Slater 1995).

References

China, W.E. and Usinger, R.L. 1949. A new genus of Tribelocephalinae from Fernando Poo. *Annali del Museo Civico di Storia Naturale de Genova*, **64**: 43-47.

Maldonado, J.C. 1996. New taxa and key to the tribes and genera in Tribelocephalinae Stål 1866 (Heteroptera: Reduviidae). *Proceedings of the Entomological Society of Washington*, **98**: 138-144.

Rédei, D. 2007. A new genus of tribelocephaline assassin bugs from Borneo (Hemiptera: Heteroptera: Reduviidae). *Zootaxa*, **1465**: 47-53.

Stål, C. 1866. *Analecta hemipterologica*. *Berliner Entomologische Zeitschrift*, **10**: 151-172.

Villiers, A. 1943. Morphologie et systématique des Tribelocephalidae africains. *Revue Française d'Entomologie*, **10**: 1-28.

Weirauch, C. 2008. Cladistic analysis of Reduviidae (Heteroptera: Cimicomorpha) based on morphological characters. *Systematic Entomology*, **33**: 229-274.

Weirauch, C. 2010. *Tribelocodia ashei*, new genus and new species of Reduviidae (Insecta: Hemiptera), has implications on character evolution in Ectrichodiinae and Tribelocephalinae. *Insect Systematics and Evolution*, **41**: 103-122.

Tribelocephala
sp., Tanzania



© Christiane Weirauch



Xenocaucus mancinii, Bioko

Opistoplatys sp.,
Indonesia



Vesciinae

[Back to Content](#)

[Back to Key](#)

Diagnosis: Vesciinae are characterized by the foretibial spur projecting beyond the tarsal insertion, the large anterior pronotal lobe, and the scape shorter than or as long as the head. Most species do not possess ocelli (except *Mirambulus* Breddin) and some genera have a frontal spine (*Chopardita* Villiers and *Vescia* Stål) and an elongate forecoxa (*Microvescia* Wygodzinsky, *Pessoaia* Costa Lima, and *Vescia*).

Taxonomic History and Distribution: The group consists of 5 genera and 20 species (Maldonado 1990). Vesciinae were established as a subfamily by Fracker and Bruner (1924) for the genus *Vescia*, which was originally described by Stål (1866) in the subfamily Reduviinae. Members of 3 genera (*Mirambulus*, *Pessoaia*, and *Vescia*) were included in Weirauch's (2008) morphological cladistic analysis of Reduviidae. *Mirambulus* and *Vescia* were recovered as sister taxa, but *Pessoaia* as the sister taxon to a large clade comprising other reduviid taxa (e.g., Reduviinae, Peiratinae, Salyavatinae), thus rendering Vesciinae polyphyletic. The analysis by Hwang and Weirauch (2012) included only one species of *Mirambulus* and did therefore not test these hypotheses. One genus (*Chopardita*) is restricted to Africa, the remaining genera are Neotropical.

Occurrence in Canada: Vesciinae do not occur in Canada.

Natural History: The natural history of Vesciinae is largely unknown. *Mirambulus morio* Breddin has been found in disturbed habitats alongside roads or in secondary forests, and *Pessoaia limai* Usinger was collected in Malaise traps and at light in tropical dry forests (Forero 2006). Brachyptery may occur in females of *Vescia* spp. (Forero 2006).



Vescia sp.,
Brazil

© Christiane Weirauch



Mirambulus sp.,
French Guiana

[Next](#)

Vesciinae (cont.)

References


Forero, D. 2006. New records of Reduviidae (Hemiptera: Heteroptera) from Colombia and other Neotropical countries. *Zootaxa*, **1107**: 1-47.

Fracker, S.D. and Bruner, S.C. 1924. Notes on some Neotropical Reduviidae. *Annals of the Entomological Society of America*, **17**: 163-174.

Stål, C. 1866. Bidrag till Reduviidernas kaennedom. Öfversigt af Kongliga Vetenskaps-Akademiens Förhandlingar, **23**: 235-302.

Weirauch, C. 2008. Cladistic analysis of Reduviidae (Heteroptera: Cimicomorpha) based on morphological characters. *Systematic Entomology*, **33**: 229-274.

[Previous](#)



Mirambulus sp., Costa Rica

© Stephen Marshall



Pessoaia sp.,
French Guiana

© Christiane Weirauch

Visayanocorinae

Diagnosis: Visayanocorinae are characterized by a foretibial spur projecting beyond the tarsal insertion, the small size, a very long second (first visible) labial segment, long second tarsomere, and the scape longer than the head. Members of this subfamily also have very smooth cuticle and lack ocelli and an antecular sulcus.

Taxonomic History and Distribution: This group consists of only 2 genera, *Carayonia* Villiers and *Wardamanocoris* Malipatil, comprising 10 species. Visayanocorinae were established by Miller (1952) for a monotypic genus from the Philippines, but it was later synonymized with the saicine genus *Carayonia* by Villiers (1958). China and Miller (1959) disagreed with Villiers' classification and retained Visayanocorinae. Subsequently, the group was either treated as a distinct subfamily (Putshkov and Putshkov 1985) or as part of the Saicinae (Maldonado 1990). Villiers (1969) provided a key to the African species of *Carayonia*. Members are found in Australia and the Old World tropics.

Occurrence in Canada: Visayanocorinae do not occur in Canada.

Natural History: Little is known on the natural history of Visayanocorinae. They have been collected using light traps, flight intercept traps, or by sieving rainforest litter (Malipatil 1990). Females of *Carayonia camerunensis* Villiers may be micropterous (Villiers 1958).

[Back to Content](#)

[Back to Key](#)



© Christiane Weirauch

Carayonia sp., Australia



[Next](#)

Visayanocorinae

References

China, W.E. and Miller, N.C.E. 1959. Check-list and keys to the families and subfamilies of the Hemiptera-Heteroptera. Bulletin of the British Museum (Natural History) (Entomology), **8**: 1-45.

Malipatil, M.B. 1990. First record of Visayanocorinae (Hemiptera: Reduviidae) from Australia with description of a new genus and two new species. Australian Journal of Entomology, **29**: 31-36.

Miller, N.C.E. 1952. Three new subfamilies of Reduviidae (Hemiptera-Heteroptera). Eos, **28**: 85-90.

Villiers, A. 1958. Hemipteres Reduviides recoltés en Angola (3 note). Publicações Culturais da Companhia dos Diamantes de Angola Lisboa, **38**: 11-46.

Villiers, A. 1969. Revision des Reduviides africains. IV. Saicinae. Bulletin de l'Institut Fondamental d'Afrique Noire, Ser. A, **31**: 1186-1247.

[Previous](#)

Carayonia sp.,
Australia



© Christiane Weirauch

